



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

INDEX TO VOLUME II.

- ABBOTT, C. C., notice of his 'A Naturalist's Rambles about Home,' 86.
- Acanthidops bairdi*, 291.
- Accipiter cooperi*, 17, 284, fuscus, 17, 174, 284, 311.
- Actodromas bairdi*, 18, 63, 273, 384.
- fuscicollis*, 63, 286.
- maculata*, 18, 286.
- minutilla*, 18, 63, 192, 268.
- Ægialites*, sp., 157.
- cantianus*, 335.
- melodus*, 37.
- melodus circumcinctus*, 285.
- mongolicus*, 240.
- semipalmatus*, 201, 241, 285.
- vociferus*, 375.
- Ægiothus canescens exilipes*, 63.
- linaria*, 16, 99, 200, 240, 279, 291.
- Æsalon columbarius*, 17, 285.
- richardsoni*, 17, 205, 285.
- Æthya americana*, 10, 18, 288.
- vallisneria*, 18, 288.
- Agelæus phœniceus*, 8, 16, 114, 142, 238, 249, 282.
- Agersborg, G. S., the birds of South-eastern Dakota, 276.
- Aix sponsa*, 10, 18, 192, 288, 369, 372.
- Alauda blakistoni*, 95.
- Albatross*, Short-tailed, 175.
- Wandering*, 387.
- Alca cristatella*, 341, 342.
- impennis*, 115.
- psittacula*, 342.
- Alcedo ispida*, 95.
- Aldrich, Charles, a belated bird, 308.
- Alle nigricans*, 211.
- Allen, J. A., sexual selection and the nesting of birds, 129; capture of escaped cage-birds, 314; a Hawk Owl at Chatham, Mass., 383; a Crested Auk on the Massachusetts-coast, 388.
- Aluco flammeus americanus*, 284.
- flammeus pratincola*, 192.
- American Ornithologists' Union, notes respecting, 223, 317, 318, 389.
- Ammodramus caudacutus nelsoni*, 216, 306.
- maritimus*, 272.
- savanarum passerinus*, 371.
- Ampelis cedrorum*, 15, 279.
- garrulus*, 15, 279.
- phœnicopterum*, 291.
- Amphispiza belli*, 16.
- belli nevadensis*, 16.
- bilineata*, 226, 228.
- Anas americana*, 158.
- boschas*, 18, 144, 158, 112, 288, 372.
- carolinensis*, 158.
- nivalis*, 207.
- obscura*, 18, 372.
- Anorthura fumigata*, 291.
- trogodytes hyemalis*, 15, 113.
- Anser albifrons gambeli*, 63, 158, 287.
- hyperboreus nivalis*, 207.
- hutchinsi*, 292.
- leucopareia*, 292.
- Anthus japonicus*, 290.
- ludovicianus*, 15, 209, 278, 290.
- pennsylvanicus*, 157, 290, 368.
- spinoletta*, 290, 368.
- spipoletta*, 290, 368.
- spraguei*, 368.
- Antrostomus carolinensis*, 220.
- vociferus*, 179.
- Aphelocoma sieberi arizonæ*, 371.
- sordida arizonæ*, 356, 371.
- woodhousei*, 16.
- Aquila chrysaëtus*, 205.
- chrysaëtus canadensis*, 10, 17, 157, 285.
- Archibuteo ferrugineus*, 17.
- lagopus sancti-johannis*, 17, 285, 295.
- Arctonetta fischeri*, 158.
- Ardea æræulea*, 272.
- egretta*, 336.
- herodias*, 10, 17, 143, 190, 287.
- Arenaria interpres*, 157.

- Asio accipitrinus*, 9, 143, 157, 189, 201, 241, 284.
 americanus, 17, 284.
 wilsonianus, 192.
Astragalinus psaltria, 16, 141.
 tristis, 16, 141, 279.
Astur atricapillus, 212, 284.
 atricapillus striatulus, 205.
Asturina nitida plagiata, 205.
Asyndesmus torquatus, 356.
 Atkins, Dr. H. A., notice of death of, 391.
Attila cozumelæ, 295.
 Auk, Crested, 388.
 Great, 115, 209, 374.
 Paroquet, 388.
 Razor-billed, 38.
Auriparus flaviceps, 229.
 Avocet, American, 285, 311.
 Bagg, Egbert, Jr., the Turkey Buzzard in Central New York, 109.
 Bain, Francis, the migration of Swallows, 216; a Blue Heron's meal, 221; winter birds of Prince Edward Island, 262.
 Baldpate, 288.
 Ball, Forrest, *Colaptes auratus* in California, 383.
 Banks, James W., nest and eggs of the Rusty Grackle (*Scolecophagus ferrugineus*), 106.
 Barrows, W. B., abnormal coloration in a caged Robin, 303; another black Robin, 303.
Bartramia longicauda, 18, 286, 315.
 Batchelder, Charles F., winter notes from New Mexico, 121, 233; *Junco annectens*—a correction, 306.
 Beckham, Charles W., notes on some of the birds of Pueblo, Colorado, 139; a white-winged Junco in Maryland, 306;
 Bell, Robert, notice of his List of Birds observed between Norway House and Forts Churchill and York, 209.
Bernicla brenta, 10, 288.
 canadensis, 18, 111, 193, 287.
 canadensis hutchinsi, 18.
 canadensis leucopareia, 293.
 nigricans, 63.
 Bicknell, Eugene P., a study of the singing of our birds, 140, 249.
 Bird, Myrtle, 99.
 Bittern, 99, 287, 374.
 Least, 101, 296.
 Blackbird, Brewer's, 282.
 Crow, 210, 253.
 Blackbird, Red-and-buff-shoulder'd, 8, 249.
 Red-winged, 210, 238, 266, 282.
 Rusty, 252, 282.
 Yellow-headed, 8, 112, 192, 282, 334.
 Blackcap, Wilson's, 332.
 Black-head, Little, 193, 288.
 Bluebird, 86, 98, 266, 277.
 Arctic, 534.
 California, 330.
 Mexican, 85, 172.
 Rocky Mountain, 8.
 Bobolink, 97, 98, 152, 189, 266, 282.
 Bob-white, 45, 315.
 Bonasa umbella, 10.
 umbella umbelloides, 17, 271.
Botaurus lentiginosus, 18, 287.
Brachyrhamphus sp., 159.
 Brant, 10, 288.
 Black, 301.
Branta canadensis hutchinsi, 158.
Brenta minima, 293.
 Brewster, William, Swainson's Warbler, 65; the Heath Hen of Massachusetts, 80; preliminary notes on some birds obtained in Arizona by Mr. F. Stephens in 1884, 84; Swainson's Warbler—an omission, 105; *Peucaea æstivalis* and its subspecies *illinoensis*, 105; Hawk Owls in New England, 109; the Eider Ducks of the New England coast, 111; the Common Cormorant in Massachusetts, 112; the Rock Ptarmigan of Newfoundland, 193; additional notes on some birds collected in Arizona and the adjoining Province of Sonora, Mexico, by Mr. F. Stephens, with a description of a new species of *Ortyx*, 196; Nelson's Sharp-tailed Finch (*Ammodramus caudacutus nelsoni*) on the Atlantic Coast, 216; the Ptarmigan of Anticosti—a correction, 220; additional notes on Swainson's Warbler (*Helinaia swainsoni*), 346; the Oyster-catcher (*Haematopus palliatus*) in Massachusetts, 384.
 Broad-bill, 193.
 Brown, Nathan Clifford, Swamp Sparrows and Yellow-rumps—a question of evidence, 307; the Great Marbled Godwit at Portland, Maine, 385.

- Browne, F. C., the Red Crossbill breeding in Eastern Massachusetts, 105; the Hawk Owl in Eastern Massachusetts, 220; another Richardson's Owl in Massachusetts, 384.
- Bubo virginianus*, 212, 284, 298.
 virginianus arcticus, 17, 21.
 virginianus saturatus,
 virginianus subarcticus, 8, 143.
- Bucephala albeola*, 337.
 islandica, 336.
- Budytes flavus*, 157.
- Buffle-head, 193.
- Bunting, Baird's, 268.
 Bay-winged, 23.
 Black-throated, 106, 192, 281, 307.
 Henslow's, 209.
 Indigo, 266, 289.
 Lark, 32, 49, 281.
 Snow, 101, 265, 279, 290.
 Varied, 198.
- Burgomaster, 37.
- Buteo borealis*, 174, 285.
 borealis calurus, 9, 17, 356.
 borealis krideri, 109.
 borealis socorroensis, 205.
 brachyurus, 204.
 cooperi, 17, 165.
 fuliginosus, 204, 371.
 lineatus alleni, 292.
 oxypterus, 371.
 swainsoni, 226, 285.
 swainsoni oxypterus, 371.
- Bryant, Walter E., the relationship of *Podiceps occidentalis* and *P. clarkii*, 313.
- Buzzard, Swainson's, 285.
 Turkey, 10, 109, 213, 285.
- CÆLIGENA clemenciæ*, 85, 199.
- Calamospiza bicolor*, 16, 32, 226, 281.
 melanocorys, 49.
- Calcarius lapponicus*, 157.
- Calipepla californica vallicola*, 372.
 squamata, 213, 226, 228.
- Calodromas*, 208.
- Calopezus*, 208.
- Calypte costæ*, 231, 309.
- Campylorhynchus brunneicapillus*, 220, 229.
- Canace obscura*, 10, 17.
- Cancroma zeledoni*, 370.
- Canvasback, 228.
- Caprimulgus minor americanus*, 178, 179.
- Caprimulgus virginianus*, 178.
 vociferus, 257, 272, 283.
- Cardellina rubrifrons*, 353.
- Cardinal, 72.
- Cardinalis cardinalis suberbus*, 344.
 saturatus, 295.
 virginianus, 15, 16, 61, 307.
- Carpodacus cassinii*, 16, 73, 333, 349.
 frontalis, 16, 141.
 frontalis mexicanus, 207.
 hæmorrhous, 207.
 mexicanus, 207.
 purpureus, 279.
- Carpodectes antoniae*, 291.
- Catbird, 55, 98, 99, 215, 266, 277, 297, 299, 300.
- Catharista atrata*, 109, 311.
- Cathartes aura*, 10, 17, 109, 143, 285, 356.
 burrovianus, 293.
 urubitinga, 293.
- Catherpes mexicanus*, 233.
 mexicanus conspersus, 15, 350.
- Cazique, Blue-throated, 85, 199.
- Ceophlæus*, 52.
 pileatus, 52.
- Centrocerus urophasianus*, 10, 17.
- Centurus carolinus*, 17, 283, 335.
 dubius, 295.
 uropygialis, 226, 228.
 leei, 295.
- Centronyx bairdi*, 16, 198, 231, 333.
- Centrophanes lapponicus*, 16, 63, 190, 280.
 ornatus, 16, 280, 333.
 pictus, 190, 280.
- Cepphus carbo*, 294.
 columba, 159, 294.
 grylle, 294.
 mandtii, 294.
 motzfeldi, 294.
- Ceratorhina moneracata*, 387.
- Cerorhina suckleyi*, 388.
- Certhia familiaris*, 291.
 familiaris mexicana, 197, 350.
 familiaris rufa, 277.
 familiaris rufa montana, 15.
 pinus, 343.
- Certhiola finschi*, 293.
 sancti-thomæ, 293.
 sundevalli, 293.
 tricolor, 291.
- Ceryle alcyon*, 9, 17, 238, 260, 283, 311.
 superciliosa stictoptera, 208.
- Chadbourne, Arthur P., *Oporornis agilis* and *Dendroica palmarum*

- palmarum* at Shelburn, New Hampshire, 104; wintering of the Swamp Sparrow in Eastern Massachusetts, 216; Swamp Sparrows and Yellow-rumps, 380.
- Chætura pelagica*, 257, 283.
yucatanica, 290.
- Chamæpelia passerina*, 229.
- Chamberlain, Montague, the nesting habits of the Cape May Warbler (*Dendræca tigrina*), 33; albino Robins (*Turdus migratorius*), 102; a white Crow (*Corvus frugivorus*), 107; the Kingbird in a new rôle, 108; non-appearance of Juncos at Montreal, 380.
- Charadrius dominicus*, 18, 285.
dominicus fulvus, 157.
- Chat, Yellow-breasted, 101, 278, 308.
- Chaulelasmus streperus*, 18, 112, 144, 288.
- Chen caerulescens*, 10, 287.
hyperboreus, 18, 287.
hyperboreus albatrus, 63, 207.
- Chewink, 150, 308.
- Chickadee, Black-capped, 265, 304.
 Carolina, 304.
 Hudsonian, 265.
 Long-tailed, 277, 330.
 Mountain, 126, 127, 330.
- Chlorostilbon caniveti*, 295.
forficatus, 295.
- Chondestes grammica*, 281.
grammica strigata, 16, 141.
- Chordeiles popetue*, 55, 178, 283.
popetue henryi, 17, 143.
virginianus, 178, 180, 257.
virginianus henryi, 180.
virginianus minor, 180.
- Chroicocephalus franklini*, 222.
- Chrysomitris pinus*, 16, 124, 141, 210, 333.
- Chuck-will's-widow, 211, 220.
- Cinclus mexicanus*, 8, 15, 140, 157, 234, 330.
- Circus cyaneus hudsonius*, 22, 63, 193.
hudsonius, 9, 17, 157, 284.
- Cistothorus stellaris*, 190, 278.
- Clangula albeola*, 18, 158, 193, 288.
glaucium, 373.
glaucium americana, 18.
islandica, 18.
- Coale, H. K., a new bird for Illinois, 109.
- Coccyzus americanus*, 17, 283, 335.
erythrophthalmus, 283.
- Cock, Sage, 10.
- Colaptes auratus*, 259, 284, 335, 383.
auratus mexicanus, 8, 143.
chrysoides, 226, 228, 302.
mexicanus, 127, 174, 302, 335, 356.
- Colinia*, 44.
- Colinus*, 44, 45.
graysoni, 45.
ridgwayi, 199, 228, 231.
virginianus, 45.
virginianus floridanus, 45.
virginianus texanus, 45.
- Columba fasciata*, 17, 174.
- Colymbus* sp., 158.
adamsi, 63.
nigricollis, 340, 341.
pacificus, 63.
septentrionalis, 63.
torquatus, 10, 18, 63, 288.
- Contopus albicollis*, 290.
borealis, 17.
depressirostris, 294.
pertinax, 356.
pileatus, 292.
richardsoni, 17, 143.
viridis, 63, 255, 369.
- Conurus carolinensis*, 17.
- Cooke, W. W., notes on the occurrence of certain birds in the Mississippi Valley, 31; Indian bird names, 222.
- Coot, American, 10, 112, 287.
- Cormorant, 210, 211.
- Cormorant, Common, 97, 112.
 Double-crested, 288.
- Corvus americanus*, 334.
corax carnivorus, 9, 16, 157, 174, 282, 371.
corax sinuatus, 371.
cryptoleucus, 16.
frugivorus, 9, 16, 107, 253, 282.
grebnitzskii, 95.
- Cory, C. B., notice of his 'Birds of Haiti and San Domingo,' 295; notice of his 'List of the Birds of the West Indies,' 319.
- Cotile riparia*, 15, 63, 279, 315.
- Coturniculus henslowi*, 190, 209, 281.
lecontei, 23, 190, 381.
mexicanus, 370.
passerinus, 63, 281, 371.
passerinus perpallidus, 16.
- Coues, Elliott, probable occurrence of *Diomedea exulans* in Florida, 387.
- Cowbird, American, 49, 107, 152, 309.

- Cowbird, Bronzed, 211.
 Crane, Corn, 296, 297.
 Crane, Sandhill, 10, 287.
 Whooping, 287.
 Creeper, Black-and-White, 112, 278.
 Brown, 98, 126, 265, 277.
 Honey, 293.
 Mexican, 197.
 Crex pratensis, 296, 297.
 Crossbill, American, 8.
 Red, 98, 100, 105, 207, 264,
 297, 333, 379.
 White-winged, 98, 264.
 Crow, Clarke's, 101, 282.
 Common, 9, 107, 213, 253,
 282, 296, 298, 300, 334, 373.
 Crymophilus, 182.
 fulicarius, 182.
 Cuckoo, Black-billed, 283.
 Yellow-billed, 283, 335.
 Cupidonia cupido, 17, 82, 285.
 pinnata, 82.
 Curlew, Eskimo, 41, 273, 287.
 Hudsonian, 10, 287.
 Long-billed, 10, 41, 273, 286.
 Cyanecula suecica, 240.
 Cyanocitta cristata, 252, 282.
 stelleri frontalis, 217.
 stelleri macrolopha, 16, 125,
 142, 174, 355.
 Cyanocorax cucullatus, 293.
 Cyanolesbia, 46, 47.
 forficata, 47.
 mocoa, 47.
 smaragdina, 47.
 Cyclorhis insularis, 295.
 Cyclorhynchus psittacula, 159, 388.
 Cygnus columbianus, 193.
 Cymbilanius lineatus fasciatus, 294.
 Cymochorea leucorrhoa, 95, 158.
 Cypselus saxatilis, 6, 17.
 Cypseloides niger borealis, 17.
 Cyrtonyx montezumæ, 46.
 DAB-CHICK, 314.
 Dafila acuta, 18, 63, 193, 201, 241,
 288, 372.
 Damon, J. L., Cowbird wintering
 in Western New York, 309.
 Davison, J. L., the first nest and
 eggs of *Eremophila alpestris*
 found in Niagara County, N. Y.,
 217.
 Dendroeca æstiva, 15, 141, 278, 331.
 auduboni, 15, 141, 331, 352.
 blackburniæ, 103, 376.
 bryanti, 370.
 bryanti castaneiceps, 370.
 cærulea, 15, 278, 369.
 cærulescens, 62, 65.
 Dendroeca castanea, 65, 272, 315.
 coronata, 15, 216, 278.
 discolor, 62, 278.
 graciæ, 15, 331, 352.
 kirtlandi, 376, 381.
 maculosa, 15, 65, 278, 376.
 nigrescens, 15, 197, 352.
 palmarum hypochrysea, 315.
 palmarum palmarum, 104.
 petechia rufivertex, 295.
 pinus, 278, 3.
 rufopileata, 291.
 striata, 61, 278.
 tigrina, 33.
 townsendi, 331.
 vieilloti, 370.
 vigorsii, 343.
 Dendrocopos immaculatus, 95.
 Diomedea albatrus, 158.
 brachyura, 175.
 exulans, 387.
 Dolichonyx oryzivorus, 16, 62, 152,
 189, 282.
 Dove, Mourning, 10, 285.
 Trinidad Ground, 358.
 Drew, Frank M., on the vertical
 range of birds in Colorado, 11.
 Dryobates, 52.
 Dryopicos, 52.
 Dryopicus, 52.
 Dryotomus, 52.
 Duck, Buffle-head, 337.
 Dusky, 336.
 Eider, 43, 111, 113, 266.
 Harlequin, 337.
 Labrador, 209.
 Lesser Scaup, 193.
 Long-tailed, 288, 385.
 Pin-tail, 193.
 Redhead, 10, 288.
 Ring-billed, 288.
 Ring-neck, 163.
 Ruddy, 288.
 Shoveller, 193, 288.
 Sprig-tail, 193.
 Summer, 193.
 Surf, 386.
 Wood, 10, 193, 288.
 Dutcher, William, Bird Notes from
 Long Island, N. Y., 36; the Can-
 ada Goose, 111.
 Dwight, J., Jr., the Ipswich Spar-
 row (*Passerculus princeps*) in
 Delaware, 105.
 Dytes auritus, 96, 289.
 auritus californicus, 18.
 nigricollis californicus, 96,
 289.

- EAGLE, Bald, 10, 192, 285.
 Golden, 10, 98, 285.
 White-headed Sea, 192.
 Ectopistes migratorius, 285.
 Egret, Great White, 110, 192, 336.
 Eider, Pacific, 386.
 Elainea cinerescens, 291.
 Elanoides forficatus, 19, 21, 192.
 Elanus leucurus, 205.
 Empidonax acadicus, 112, 256.
 atriceps, 291.
 difficilis, 17.
 flaviventris, 65, 256.
 fulvifrons, 207.
 fulvifrons pallescens, 207.
 fulvifrons rubicundus, 207.
 gracilis, 295.
 hammondi, 17, 143, 283.
 minimus, 17, 256, 283.
 obscurus, 17.
 pusillus, 17.
 trailli, 256.
 viridescens, 291.
 Engyptila gaumeri, 290.
 Eniconetta stelleri, 158.
 Ephialtes, 183.
 Eremophila alpestris, 189, 217.
 alpestris crysolæma, 17, 239.
 alpestris leucolæma, 16, 142.
 Ereunetes occidentalis, 384, 385.
 pusillus, 18, 63, 143, 201, 241, 286.
 pusillus occidentalis, 110.
 Erismatura rubida, 144, 288.
 Euetheia, 48.
 bicolor, 48.
 olivacea intermedia, 395.
 zena, 48.
 Eugenes fulgens, 199.
 FALCO albigularis, 205.
 arcticus, 188.
 candicans, 90, 187.
 candicans β islandicus, 188.
 candicans gyrfalco, 90, 188.
 communis, 267.
 fuscus, 188.
 grœnlandicus, 187, 188.
 gyrfalco, 90, 187, 188.
 gyrfalco candicans, 90, 188.
 gyrfalco grœnlandicus, 188.
 gyrfalco islandicus, 188.
 holbœlli, 187.
 islandicus, 188, 315.
 islandicus candicans, 188.
 islandus, 185, 187, 188.
 islandus β albus, 187.
 islandus γ maculatus, 187.
 islandus fuscus, 188.
 Falco peregrinus nævius, 9, 17, 212, 285.
 peregrinus pealei, 157.
 rusticolus, 185, 188.
 rusticolus gyrfalco, 187.
 rusticolus obsoletus, 187.
 FALCON, Peregrine, 267.
 Prairie, 285.
 Finch, Bachman's, 106.
 Cassin's, 173, 333.
 Grass, 280.
 Lark, 281.
 Lazuli, 281.
 Nelson's Sharp-tailed, 306.
 Painted, 72, 298.
 Pine, 124, 239, 333.
 Purple, 263, 279.
 Sea-side, 272, 301.
 Sharp-tailed, 216, 301, 306.
 Fisher, A. K., Wood Ibis (*Tantalus loculator*) in Eastern New York, 222; capture of *Ammodramus caudacutus nelsoni* in the Lower Hudson Valley, New York, 306; capture of two more specimens of *Helminthophila leucobronchialis* at Sing Sing, New York, 378; evidence concerning the interbreeding of *Helminthophila chrysoptera* and *H. pinus*, 378.
 Flicker, Golden-winged, 284.
 Flycatcher, Acadian, 112.
 Arizona Crested, 199.
 Ash-throated, 98.
 Black-crested, 242.
 Great-crested, 254.
 Hammond's, 283.
 Least, 256, 283.
 Olivaceous Crested, 199.
 Small Green-crested, 256.
 Swallow-tailed, 218.
 Traill's, 256.
 Wilson's, 330.
 Yellow-bellied, 256.
 Fratercula corniculata, 159.
 Frost, Edwin Brant, the Logger-head Shrike in New Hampshire, 379.
 Fulica americana, 10, 18, 112, 287.
 caribæa, 292.
 spinosa, 337.
 Fringilla æstivalis, 106.
 bachmani, 105.
 bicolor, 48, 49.
 mexicana, 207.
 zena, 48.
 Fuligula affinis, 193.
 collaris, 193.

- Fuligula ferina americana*, 193, 372.
Fulix, *affinis*, 18, 288.
 collaris, 18, 158, 288.
 marila, 18, 158.
Fulmarus glacialis glupischa, 158.
 GADWALL, 112, 288.
Galeoscoptes carolinensis, 15, 140, 277.
Gallinago media wilsoni, 18.
 wilsoni 286, 336.
Gallinula galeata, 18, 193, 210.
Gallinule, Common, 193.
 Florida, 193.
 Gannet, 98.
Gazetta candidissima, 17, 299.
 Gault, B. T., nest and eggs of *Calypte costae*, 309.
Geococcyx californianus, 17, 143, 375.
Geothlypis bairdi, 294.
 macgillivrayi, 15, 141, 332.
 philadelphia, 65, 278.
 speciosa, 368.
 trichas, 15, 62, 65, 278, 367.
 trichas occidentalis, 141.
Glaucidium phalænoides, 229, 231.
 Gnat-catcher. Blue-gray, 272, 334.
 Godwit, Hudsonian, 286.
 Marbled, 286, 385.
 Golden-eye, 266.
 Barrows, 336.
 Goldfinch, 264, 279.
 Pine, 264.
 Goodale, Joseph L., the Wood Thrush in Maine, 215; the Yellow-rumped Warbler wintering in Maine, 216.
 Goss, N. S., observations on *Elanoides forficatus* and *Ictinia subcærulea* in Kansas, 19; rare summer residents in Kansas, 113; *Cyanocitta stelleri frontalis* nesting in holes in trees, 217; Wilson's Plover in Nova Scotia, 221; *Rissa tridactyla kotzebuei* in Washington Territory, 222; the Black-capped Vireo and Nonpareil in Southwestern Kansas, 274; early accidental occurrence of *Catharista atrata* and *Tantalus loculator* in Kansas, 385; the Little Yellow Rail (*Porzana noveboracensis*) in Kansas, 385; the Thick-billed Grebe (*Podilymbus podiceps*) breeding in Kansas, 388.
 Goose, American White-fronted, 287.
 Canada, 42, 111, 193, 214, 287.
 Blue-winged, 10, 287.
 Goose, Common Wild, 193.
 Snow, 287.
 Goshawk, 266, 284.
 Grackle, Boat-tailed, 100.
 Bronzed, 282.
 Purple, 189.
 Rusty, 106, 282, 373.
Granatellus sallæi boucardi, 273.
 Grebe, American Eared, 96, 289.
 Carolina, 96.
 Clarke's 313.
 Eared, 340.
 European Crested, 93.
 Horned, 96, 193, 289, 314, 340.
 Red-necked 314.
 Thick-billed, 289, 388.
 Western, 313, 314.
 Greenlet, 117.
 Grosbeak, Black-headed, 100, 112, 192, 281.
 Blue, 112, 211.
 Cardinal, 97, 98, 151, 307, 343.
 Evening, 174, 279, 296, 334.
 Pine, 264.
 Rose-breasted, 100, 151, 266, 281, 299.
 Grouse, Dusky, 10.
 Pinnated, 80.
 Ruffed, 10, 40, 270, 300.
 Sharp-tailed, 285.
 Willow, 201.
Grus americanus canadensis, 10, 18, 63, 144, 157, 201, 287.
 Guillemot, Brünnich's, 38.
Guiraca cærulea, 16, 112, 142, 211.
 Gull, Black-backed, 266.
 Bonaparte, 266, 374.
 Burgomaster 37.
 Franklin's 289.
 Glaucous, 37, 337.
 Herring, 266, 288.
 Kittiwake, 266.
 Pacific Kittiwake, 222.
 Ring-billed, 288.
 Sabine's, 374, 375.
 Gurney, J. H., notice of his 'List of the Diurnal Birds of Prey,' 203.
Gymnocitta cyanocephala, 9, 13, 16.
 Gyrfalcon, 187.
 Gray, 187, 188.
 Iceland, 284.
 Labrador 187.
 White, 187.
HÆMATOPUS bachmani, 157.
 *palliatu*s, 384.
Haliaëtus leucocephalus, 10, 16, 143, 157, 192, 285.

- Hardy, Manly, late occurrence of the Phoebe (*Sayornis fuscus*) at Brewer, Maine, 108.
- Harelda glacialis, 63, 288.
hyemalis, 158, 385.
- Harporhynchus cinereus bendirei, 15.
curvirostris palmeri, 196, 228.
guttatus, 294.
lecontei, 196, 229, 230, 231.
melanostoma, 294.
rufus, 15, 140, 277, 298.
- Harrier, Common, 22, 192.
- Hawk, Duck, 9, 209, 285.
Fish, 9, 192, 210, 212, 285.
Florida Red-shouldered, 292.
Marsh, 9, 100, 192, 285.
Pigeon, 100, 285.
Red-tailed, 174, 285.
Rough-legged, 285, 295.
Sharp-skinned, 99, 174, 284, 311.
Sparrow, 100, 285.
Western Red-tailed, 9.
- Hayward, R., curious food for the Kingfisher (*Ceryle alcyon*), 311.
- Helinaia, 76.
swainsoni, 76, 105, 346, 377.
- Helmintherus swainsoni, 301.
vermivorus, 62, 103, 304.
- Helminthophaga celata, 15, 40, 211, 278, 331.
peregrina, 15.
ruficapilla, 15.
virginiae, 15, 331.
- Helminthophila celata lutescens, 352, 366.
chrysoptera, 102, 304, 359, 361, 363, 378.
gunni, 362.
lawrencei, 359, 361, 363.
leucobronchialis, 304, 359, 378, 361.
peregrina, 65.
pinus, 359, 361, 363, 378.
virginiae, 352.
- Helonæa swainsoni, 62, 104.
- Hen, Heath, 80, 82.
Prairie, 81, 82, 285.
- Henshaw, H. W., the Gulls of the Californian Coast, 231; list of birds observed in summer and fall on the upper Pecos River, 326; a new bird to Massachusetts, 384; the Baird's Sandpiper (*Actodromas bairdi*) at Locust Grove, New York, 384; the bill of the Horned Puffin (*Ceratorhina monocerata*), 387.
- Hierodias alba egretta, 96, 299.
egretta, 110, 192.
- Heron, Black-crowned Night, 10, 192.
Great Blue, 10, 190, 221, 287.
Green, 287.
Little Blue, 272.
Night, 97, 287.
White, 96, 100, 192.
- Hesperiphona vespertina, 15, 174, 279, 334, 349.
- Heteractitis incanus, 157.
- Hierofalco gyrfalco, 204.
gyrfalco candicans, 205.
gyrfalco islandus, 205, 284.
gyrfalco obsoletus, 205.
gyrfalco sacer, 63.
holbœlli, 205.
mexicanus polyagrus, 17, 205, 285.
- Himantopus mexicanus, 18.
- Hirundo erythrogaster, 15, 141, 278, 366.
rustica, 366.
rustica erythrogastra, 366.
- Histrionicus histrionicus, 158, 337, minutus, 18.
- Hoffman, W. J., bird names of the Selish, Pah-Uta, and Shoshoni Indians, 7.
- Holterhoff, Godfrey, Jr., the Glossy Ibis and Avocet at San Diego, Cal., 311.
- Hoy, Dr. P. R., nest and eggs of the Golden-winged Warbler (*Helminthophila chrysoptera*), 102.
- Hummingbird, Black-chinned, 310.
Costa, 310.
Refulgent, 199.
Ruby-throated, 9, 218, 256, 283, 299, 300.
- Hydrochelidon lariformis, 211.
lariformis surinamensis, 18, 112.
surinamensis, 289.
- Hylocichla fuscescens, 277.
fuscescens salicicola, 15.
mustelina, 278, 369.
unalasca, 330.
unalashkæ aububoni, 15, 330.
ustulata swainsoni, 15, 140, 277, 330.
- Hylotoma, 51, 52.
- Hylotomus, 51, 52.
pileatus, 283.
- IBIS, Glossy, 311.
White, 287.
Wood, 213, 221, 311.
- Icteria virens, 278.
virens longicauda, 15, 141.

- Icterus bullocki*, 16, 142, 282.
 cucullatus, 159, 292.
 cucullatus igneus, 292.
 cucullatus nelsoni, 292.
 curasoënsis, 291.
 galbula, 16, 55, 58, 251, 282, 298.
 parisorum, 1, 355.
 spurius, 16, 251, 282.
Ictinia subcærulea, 19.
 Indigo-bird, 151.
 Ingersoll, Ernest, notice of his 'Country Cousins; Short Studies in the Natural History of the United States,' 91.
Ionornis martinica, 63.
Iridoprocne bicolor, 189.
 JACANA, Brazilian, 337.
 Jay, Blue, 100, 101, 253, 265, 282.
 Blue-fronted, 217.
 Canada, 99, 107, 265, 335.
 Long-crested, 125.
 Piñon, 9.
 Steller's, 174.
 Jeffries, Wm. A., the Common Cormorant off Boston Harbor, 112.
 Jer-Falcon, Brown, 90.
 White, 90.
 Jordan, David Starr, notice of his 'Manual of the Vertebrates of the Northern United States,' 208.
Junco, 115, 116, 117, 380.
 Mexican, 198.
 Slate-colored, 146.
 White-winged, 306.
Junco aikenii, 16, 32.
 annectens, 16, 306.
 caniceps, 16, 123, 124, 141.
 cinereus, 174, 198, 363, 364.
 cinereus caniceps, 174, 355.
 cinereus dorsalis, 354, 364.
 cinereus palliatus, 354, 364.
 hyemalis, 16, 63, 146, 280, 306.
 oregonus, 16, 123, 124, 141, 174.
 Kells, Wm. L., foster parents of the Cowbird, 106.
 Killdeer, 285.
 Kingbird, 108, 253, 283.
 Western, 283.
 Kingfisher, 9, 95, 238, 260, 283, 311.
 Kinglet, Golden-crested, 265, 277.
 Ruby-crowned, 93, 98, 99, 173, 277, 330.
 Kitchen, J. M. W., on the function of the inferior larynx in birds, 24.
 Kite, Mississippi, 21.
 Kite, Swallow-tailed, 19, 192, 284, 347.
 Kittiwake, 266.
 Pacific, 22.
 Knot, 312, 375.
LAGOPUS albus, 63, 221.
 albus alleni, 201.
 hyperborea, 95.
 islandorum, 95.
 lucurus, 14, 17, 95.
 muta, 94.
 muta vulgaris, 95.
 ridgwayi, 95.
 rupestris, 63, 95, 157, 194, 221.
 ruprestris atkhensis, 95.
 rupestris nelsoni, 95.
 rupestris reinhardti, 95, 195.
 welchi, 194.
Lampornis thalassinus, 295.
Lampronetta fischeri, 63, 201.
 Langille, J. H., notice of his 'Our Birds in their Haunts,' etc., 91.
 Lantz, D. E., the song of *Cardinalis virginianus*, 307.
Lanio melanopygius, 291.
Lanius borealis, 15, 212, 279.
 ludovicianus, 210, 379.
 ludovicianus excubitorides, 15, 279, 332.
Lanivireo solitarius plumbeus, 15, 141.
 Lark, Horned, 189, 217, 239, 282, 296.
 Meadow, 250, 266.
 Mexican Meadow, 198.
 Shore, 100.
 Western Meadow, 8, 282.
Larus argentatus smithsonianus, 232, 288.
 brachyrhynchus, 232.
californicus, 232.
 canus, 232.
 delawarensis, 18, 232, 288.
 franklini, 288.
 glaucescens, 158.
 glaucus, 37, 63.
 occidentalis, 232.
 philadelphæ, 232.
 Lawrence, G. N., notice of his paper describing new species of American birds, 290; descriptions of new species of the family Columbidae, 337.
Leucosticte atrata, 16.
 griseinucha, 157.
 tephrocotis, 16.
 tephrocotis littoralis, 16.
 Ligea, 295.

- Limosa fedoa*, 18, 286.
 fœda, 385
 hæmastica, 286.
 lapponica novæ-zelandiæ, 63, 201.
Lobipes hyperboreus, 18, 63, 192, 315.
Lomvia arra, 63.
 arra bruennichi, 38.
 Long-beak, Greater, 273.
 Longspur, Chestnut-collared, 280, 333.
 Lapland, 23, 190, 280, 334.
 Painted, 190.
 Smith's, 280.
 Loomis, Leverett M., supplementary notes on the ornithology of Chester County, S.C., 168.
 Loon, 289.
Lophodytes cucullatus, 18, 144, 288.
Lohophanes bicolor, 277.
 inornatus griseus, 15.
 wollweberi, 349.
Lophortyx californica, 10, 247, 372.
 californicus brunnescens, 208, 372.
 gambeli, 131, 247, 301.
 gambeli × *californica*, 247.
Loxia albigentris, 207.
 americana, 379.
 curvirostra, 207.
 curvirostra americana, 8, 16, 105, 207, 212, 333.
 curvirostra bendirei, 128, 207.
 curvirostra mexicana, 207, 372.
 curvirostra stricklandi, 372.
 leucoptera, 16, 210.
 pytiopsittacus, 207.
Lunda cirrhata, 159, 388.
 M'CALLUM, G. A., albinism, 113.
 McKinley, James, field notes from Pictou County, Nova Scotia, 39.
Macrorhamphus griseus, 18, 374.
 griseus scolopaceus, 37, 63, 273, 286.
Magpie, Black-billed, 9, 282.
Mallard, 100, 112, 289, 336.
Mareca americana, 18, 288.
 penelope, 373.
Martin, Purple, 55, 57, 98, 279, 299, 300.
Megascops, 184.
 asio, 184.
 asio bendirei, 184.
 asio floridanus, 184.
 asio kennicotti, 184.
 asio macalli, 184.
 asio maxwelliæ, 184.
Megascops flammeolus, 184.
 trichopsis, 184.
Melanerpes erythrocephalus, 17, 63, 143, 259, 283.
 formicivorus bairdi, 8, 174, 356.
 torquatus, 17, 143.
Melanetta fusca, 291.
Meleagris gallopavo, 10.
 gallopavo americana, 17, 285.
Melopelia leucoptera, 17, 229.
Melospiza cinerea, 157.
 fasciata, 95, 147, 280.
 fasciata fallax, 16, 141.
 fasciata montana, 124, 236,
 georgiana, 372,
 lincolni, 16, 65, 142, 280, 354.
 palustris, 149, 189, 216, 280, 372, 380.
Merganser, 266.
 Hooded, 112, 193.
Merganser serrator, 158.
Mergus cucullatus, 193, 372.
 merganser, 372.
 mergus americanus, 18, 144, 288.
 serrator, 201, 241, 372.
 Merlin, Richardson's, 285.
 Merriam, C. Hart, preliminary report of the Committee on Bird Migration, 53; notice of his paper on the Birds of the Bermudas, 95; nest and eggs of the Blackburnian Warbler, 103; Swainson's Warbler off Southern Florida, 104; a remarkable migration of Canada Jays, 107; the White Pelican on Lake Ontario, 111; third addendum to list of birds ascertained to occur within ten miles from Point des Montes, Province of Quebec, Canada; based chiefly upon notes of Napoleon Comeau, 113; Committee on the Migration and Geographical Distribution of North American Birds; Circular for 1885, 117; change of color in the wing-feathers of the Willow Grouse, 201; probable breeding of the Wheatear (*Saxicola ananthe*) on the northern shore of the Gulf of St. Lawrence, 305; eggs of the Knot (*Tringa canutus*) found at last, 312; fourth addendum to list of birds ascertained to occur within ten miles of Point de Monts, province of Quebec, Canada; based chiefly upon the notes of Napo-

- leon A. Comeau, 315; Kirtland's Warbler from the Straits of Mackinac, 376; odd nesting habits of the Blue Yellow-backed Warbler in Missouri, 377; Swainson's Warbler in Jamaica, 377; an albino Surf Duck (*Edemia perspicillata*), 372.
 Merrill, J. C., republication in 'The Auk' of Descriptions of New North American birds,—a suggestion, 389.
Merula merula, 129.
 migratoria, 55, 277, 303, 304.
 migratoria propinqua, 15, 140, 330.
 torquata, 129.
Micrathene whitneyi, 231.
Micrologia, 295.
Micropalama himantopus, 273, 286.
Milvulus forficatus, 218.
Mimocichla ardesiaca, 295.
Mimus carolinensis, 55, 215, 297.
 gilvus rostratus, 291.
 polyglottus, 15.
 Minor Ornithological Publications, 96, 209, 295, 372.
Mitrephorus pallescens, 207.
Mniotilta varia, 15, 62, 278.
 Mockingbird, 86, 98.
Molothrus æneus, 211.
 ater, 16, 49, 50, 106, 142, 282, 309.
 pecoris, 49, 50, 152.
Motacilla ocularis, 157.
 Murdoch, John, notes on some species of birds attributed to Point Barrow, Alaska, 200.
Muscicapa atra, 51.
 carolinensis fusca, 51.
 fulvifrons, 207.
 fusca, 51.
 nunciola, 51.
 phœbe, 51.
Myiadestes townsendi, 15, 128, 140, 331.
Myiarchus crinitus, 254.
 cinerascens, 17, 98.
 lawrencei, 208.
 lawrencei olivaceus, 198, 208.
 mexicanus magister, 198, 207.
 platyrhynchus, 295.
Myiodioctes canadensis, 65.
 minutus, 372.
 pusillus, 278, 330, 332.
NAUCLERUS forficatus, 284.
 Nelson, E. W., counter-'notes on some species of birds attributed to Point Barrow, Alaska,' 239.
Neocorys spraguei, 269.
 Nests and eggs of the birds of Ohio, notice of work on, 289.
Nettion carolinensis, 18, 144, 201, 241, 288.
 Newton, Alfred, notice of his article 'Ornithology,' 205.
Nighthawk, 55, 178, 283.
 Cuban, 182.
 Western, 182, 283.
 Night-jar, 257.
Numenius borealis, 63, 273, 287.
 hudsonicus, 10, 143, 201, 241, 287.
 longirostris, 10, 18, 208, 273, 286.
 Nutcracker, Clarke's, 282.
 Nuthatch, Pigmy, 126, 172, 330.
 Red-bellied, 265, 277.
 Slender-billed, 126, 127, 172, 277, 330.
 White-bellied, 93, 98, 265.
 Nuttall Ornithological Club, 116, 224, 390.
 Nutting, Charles C., notice of his paper on a collection of Nicaraguan birds, 293.
Nyctala acadica, 272, 284.
 tengmalmi richardsoni, 384.
Nyctea nyctea, 157.
 scandiaca, 37, 63, 284.
Nyctherodius violaceus, 18.
Nyctiardea gardeni, 299.
 grisea nævia, 10, 192, 287.
Nymphicus novæ-hollandiæ, 314.
OCEANODROMA furcata, 158.
Ochthodromus wilsonius, 221, 273.
Edemia americana, 18, 158.
 perspicillata, 158, 386.
Ednienemus dominicensis, 110.
Estrelata defilippiana, 292.
 fisheri, 292.
 Old-squaw, 266.
Olor americanus, 10.
 buccinator, 287.
 columbianus, 157.
Onychotes gruberi, 293.
 solitarius, 293.
Oreoscoptes montanus, 8, 13, 15, 140.
Oriolus ater, 49.
 niger, 50.
Oriole, Baltimore, 55, 58, 86, 101, 251, 282, 298, 290.
 Black, 49.
 Bullock's, 282.
 Hooded, 159.
 Orchard, 251, 282.
 Scotts, 1.
 Ortolan, 190.

- Ortygia, 45.
 Ortygis, 45.
 Ortyx, 45.
 Ortyx virginiana, 14, 213, 285, 296,
 297, 299, 315.
 douglasii, 46.
 elegans, 46.
 graysoni, 227, 297.
 massena, 46.
 montezumæ, 46.
 picta, 46.
 Oryzoborus nuttingi, 294.
 Osgood, Fletcher, the Chuck-will's-
 widow (*Antrostomus carolinensis*) in
 Massachusetts, 220.
 Osprey, 192.
 Oxyechus vociferus, 18, 143, 285.
 Otocoris alpestris, 282.
 Ouzel, Water, 8, 234, 330.
 Owl, Acadian, 266.
 American Long-eared, 9.
 Barn, 192, 210, 284.
 Barred, 100, 189, 266, 284.
 Burrowing, 9, 284.
 Great Horned, 99, 266, 284.
 Hawk, 31, 108, 220, 383.
 Little Screech, 9.
 Long-eared, 99, 192, 284.
 Marsh, 189.
 Monkey-faced, 101.
 Richardson's, 384.
 Saw-whet, 272, 284.
 Screech, 101, 260, 284, 296.
 Short-eared, 87, 99, 100, 189,
 284.
 Snowy, 37, 366, 284.
 Western Horned, 9.
 White, 192.
 Oyster-catcher, 384.
 PAGOPHILA eburnea, 63.
 Palmer, William, abundance of
 Parus atricapillus near Wash-
 ington, 304.
 Pandion haliaëtus, 192.
 haliaëtus carolinensis, 285.
 solitarius, 293.
 Paroquet, Cockatoo, 314.
 Parra chavaria, 338.
 dominica, 338, 339.
 jacana, 339, 338.
 senegalla, 338, 339.
 variabilis, 338.
 Parula americana, 15, 61, 62,
 377.
 nigrilora, 368.
 pitiayumi, 368.
 Parus atricapillus, 304.
 atricapillus septentrionalis,
 15, 277.
 Parus atricapillus turneri, 207, 331.
 carolinensis, 304.
 cinctus, 372, 373.
 cinctus obtectus, 372.
 montanus, 15, 126, 127, 140,
 331.
 Passer domesticus, 190, 300.
 Passerculus alaudinus, 292.
 anthinus, 212.
 bairdi, 268.
 beldingi, 292.
 princeps, 36, 105.
 rostratus, 198, 230.
 sandwichensis, 157.
 sandwichensis alaudinus, 16,
 201, 241.
 sandwichensis bryanti, 292.
 sandwichensis savanna, 16,
 272, 280.
 savanna, 65, 376.
 Passerella iliaca, 150, 281.
 iliaca schistacea, 16.
 Passerina amœna, 16, 142, 281.
 ciris, 276.
 cyanea, 16, 151, 289.
 versicolor, 198.
 Pedicætes phasianellus campes-
 tris, 208.
 phasianellus columbianus,
 285.
 Pelagadroma marina, 386.
 Pelecanus erythrorhynchus, 18, 111,
 144, 288.
 Pelican, Frigate, 113.
 White, 111, 288.
 Pelidna alpina americana, 18, 63.
 subarquata, 63.
 Pelionetta perspicillata, 95.
 Perisoreus canadensis, 335.
 Perissoglossa tigrina, 32.
 Petrel, Stormy, 99.
 White-faced Stormy, 386.
 Petrochelidon lunifrons, 15, 279,
 366.
 pyrrhonota, 366.
 Peucea æstivalis, 106.
 æstivalis bachmani, 106.
 æstivalis illinoensis, 105, 106,
 299.
 arizonæ, 226, 370.
 botteri, 370.
 carpalis, 228.
 cassini, 226.
 mexicana, 370.
 ruficeps boucardi, 354.
 Peucedramus olivaceus, 172, 349,
 352.
 Pewee, 51.
 Wood, 255.

- Phænicothraupis carmioli*, 291.
Phainopepla nitens, 228, 242.
Phalacrocorax carbo, 97, 112, 210.
 dilophus, 112, 288.
 dilophus albociliatus, 208.
 dilophus cinnatus, 158.
 pelagicus, 158.
Phalacroptilus nuttalli, 9, 17, 143,
 283, 356, 382.
Phalarope, Northern, 183, 192.
 Red, 183.
 Wilson's, 183, 273, 286.
Phalaropus fulicarius, 63.
 lobatus, 157, 183.
 tricolor, 183.
Philacte canagica, 158.
Philohela minor, 261, 286.
Philomachus pugnax, 336.
Philotrix, 45.
Phloeotomus, 52.
Phœbe, 51, 98, 108, 154, 283.
Phonipara, 48.
Phylloscopus borealis, 373.
Pica camtschatica, 95.
 rustica hudsonica, 9, 16, 142,
 282.
Picoides arcticus, 210.
 tridactylis dorsalis, 17.
Picicorvus columbianus, 16, 282.
Picus insularis, 115.
 nuttalli, 213.
 pileatus, 52.
 pubescens, 257, 283.
 pubescens gairdneri, 17, 126,
 143.
 stricklandi, 199.
 villosus, 100, 115, 373.
 villosus harrisi, 9, 17, 127,
 174, 283, 356.
Pigeon, Band-tailed, 174.
 Passenger, 285.
 Wild, 40.
Pinicola enucleator, 15.
Pintail, 288.
Pittasoma michleri zeledoni, 291.
Pipilo aberti, 16.
 chlorurus, 16, 142.
 erythrophthalmus, 150, 282,
 308, 369.
 fuscus mesoleucus, 16, 228,
 237.
 maculatus arcticus, 142, 282.
 maculatus arcticus megalonyx, 16.
 maculatus megalonyx, 128,
 355.
Plautus, 183.
Plegadis guarauna, 18.
Plectrophenax hyperboreus, 291.
 navalis, 157.
Plectrophanes lapponicus, 23, 334.
 navalis, 16, 23, 63, 279.
 ornatus, 23.
 pictus, 23.
Plover, Belted Piping, 285.
 Black-bellied, 32, 41, 285.
 Field, 286.
 Golden, 41, 285.
 Killdeer, 285, 375.
 Piping, 37.
 Semipalmated, 285.
 Snowy, 335.
 Upland, 212, 297.
 Wilson's, 221, 273.
Podasocys montanus, 18, 143.
Podiceps auritus, 340.
 californicus, 340.
 clarkii, 313.
 cornutus, 193, 314.
 griseigena holbœlli, 314.
 holbœlli, 18.
 occidentalis, 313, 314.
Podilymbus podiceps, 18, 96, 289,
 314, 388.
Pochard, American, 193.
Poliophtila cœrulea, 15, 272, 334, 349,
 plumbea, 228.
Polysticte stelleri, 63.
Poœcetes gramineus, 23, 280, 315.
 gramineus confinis, 16, 141.
Poor-will, 9, 283, 382.
Porzana carolina, 18, 62, 190, 209,
 287.
 jamaicensis, 299.
 leucogastra, 294.
 noveboracensis, 110, 274, 385.
Pratincola rubicola, 129.
Procellaria hypoleuca, 387.
 marina, 387.
Progne subis, 15, 55, 57, 228, 279,
 299.
Protonotaria citrea, 32.
Psaltiriparus grindæ, 208.
 minimus californicus, 207.
 plumbeus, 15.
Pseudogryphus californianus, 167.
Ptarmigan, Rock, 193, 222.
 Willow, 221.
Puffin, Horned, 387.
Pyranga æstiva cooperi, 15, 198,
 372.
 hepatica, 197.
 ludoviciana, 8, 15, 141, 333,
 354.
 rubra, 95, 278.
 rubra cooperi, 372.

- Pyrrhuloxia sinuata*, 228.
QUAIL, 285, 296.
 Arizona, 296, 299, 301.
 California, 10, 295.
 Florida, 45.
 Grayson's, 45, 297.
 Masked, 199.
 Massena, 46.
 Migratory, 301.
 Mountain, 300.
 Texan, 45.
Querquedula carolinensis, 18, 144, 193, 372.
 cyanoptera, 112.
 discors, 18, 144, 288, 372.
Quiscalus, 43.
 purpureus, 189.
 purpureus æneus, 16, 282.
RAIL, Carolina, 190.
 Clapper, 98, 274.
 Red-breasted, 287.
 Sora, 98, 287.
 Virginia, 96, 98, 192, 287.
 Yellow, 274, 385.
Rallus elegans, 287.
 longirostris crepitans 274.
 virginianus, 18, 144, 192, 287.
 Raven, American, 9, 174, 282.
Recurvirostra americana, 18, 144, 210, 211, 286.
 Redbird, Cardinal, 300.
 Summer, 98.
 Red-head, 193, 288.
 Redpoll, Lesser, 99, 264, 279.
 Redstart, American, 98, 278.
Regulus calendula, 14, 15, 173, 277, 311.
 japonicus, 291.
 satrapa, 15, 277, 301.
 Ricker, C. B., a specimen of *Helminthophila leucobronchialis* in New Jersey, 378.
 Ridgway, Robert, on *Buteo harlani* (Aud.) and *B. cooperi* (Cass.), 165; remarks on the Californian Vulture (*Pseudogryphus californianus*), 167; note on *Sarcorhamphus æquatorialis* Sharpe, 169; notice of his paper on the American Red Crossbills, 206; notices of his papers on various American birds, 207, 290, 370; do. of his paper on the birds of Cozumel Island, 294; description of a new Cardinal Grosbeak from Arizona, 343; *Helminthophila leucobronchialis*, 359; on *Funco cinereus* (Swain.) and its geographical races, 363; notice of his paper on the type specimen of *Buteo oxypterus*, 371; do. emended names of North American birds, 371; a Petrel new to North America, 386; announcement of his nomenclature of Colors and Compendium, etc.
 Ridgway Ornithological Club, 116, 319.
Rhodostethia rosea, 63.
Rhyacophilus solitarius, 18.
Rhynchofalco fusco-cærulescens, 405.
Rhynchophanes maccowni, 16.
Rhynchops nigra, 193, 299.
Rissa brevirostris, 158.
 tridactyla kotzbuëi, 222, 213.
 tridactyla pollicaris, 158.
 Rives, Wm. C., Jr., M. D., nesting of the Worm-eating Warbler (*Helminthophila vermivorus*) in Virginia, 103; notice of his 'The Birds of Newport,' 208.
 Robin 55, 102, 211, 277, 300, 303, 304.
 Western, 330.
 Ruff, 336.
 SAGE, John H., return of Robins to the same nesting place, 304; another example of *Helminthophila leucobronchialis* from Connecticut, 304; nesting of the Worm-eating Warbler (*Helminthophila vermivorus*) in Southern Connecticut, 305; occurrence of the Sharp-shinned Hawk in New Hampshire in winter, 311.
Salpinctes obsoletus, 15, 234, 277.
 Sandpiper, Baird's, 273, 384.
 Bartram's, 375.
 Bonaparte's, 286.
 Buff-breasted, 286.
 Least, 192, 286.
 Pectoral, 286, 374.
 Semipalmated, 286.
 Solitary, 286.
 Spotted, 286.
 Stilt, 273, 287.
 Western Semipalmated, 110, 384, 385.
Sarcogrammus cinereus, 339.
 goënsis, 339.
 indicus, 339.
 inornatus, 339.
Sarcorhamphus æquatorialis, 169.
 gryphus, 168, 269, 171.
 Saunders, W. E., the Black-throated Bunting, Yellow-breasted Chat, and Connecticut Warbler in Ontario, 307.

- Saxicola œnanthe*, 15, 113, 305.
Sayornis, 51.
 fusca, 108, 283.
 phœbe, 51, 254, 369.
 sayi, 17.
Scardafella inca, 228.
Scolecophagus cyanocephalus, 13,
 142, 282.
 ferrugineus, 106, 252, 282, 373.
Scops, 183.
 asio, 9, 184, 260, 284.
 asio maxwelliæ, 17.
 flammeolus, 17.
 indicus, 184.
Scopus, 184.
Scott, W. E. D., on the breeding
 habits of some Arizona birds; I,
 Icterus parisorum, 1; II, *Icterus*
 cucullatus, 159; III, *Phainopepla*
 nitens, 242; IV, *Vireo vicinior*,
 321; winter mountain notes from
 Southern Arizona, 172; early
 spring notes from the mountains
 of Southern Arizona, 348.
Scott, W. L., the Great White Egret
 and the Yellow Rail in Ottawa,
 Canada, 110; on the names 'Snow-
 bird' and Winter Wren, 114.
Seeböhm, Henry, notice of his 'A
 History of British Birds,' etc., 88.
Seiæphorus flammula, 291.
 platycercus, 17.
 rufus, 17, 356.
 torridus, 291.
Setophaga picta, 353.
 ruticilla, 15, 61, 62, 98, 141,
 278.
Seton, Ernest E. T., Manitoban
 Notes, 21; the Swallow-tailed Fly-
 catcher in Manitoba and at York
 Factory, 218; notes on Manito-
 ban birds, 267; nest and eggs of
 the Philadelphia Vireo, 305; the
 Western Grebe in Manitoba, 314;
 the popular names of birds, 316;
 interesting records from Toronto,
 Canada, 334.
Sheldrake, American, 288.
 Hooded, 288.
Shoveller, 112, 193, 288.
Shrike, Great Northern, 238, 278.
 Loggerhead, 379.
 White-rumped, 279, 332.
Shufeldt, R. W., notes of his me-
 moir on the osteology of *Ceryle*
 alcyon, 95; do. on the avian patel-
 la, 96; variations in the form of
 the beak, that take place during
 its growth, in the Short-tailed Al-
 batross (*Diomedea brachyura*),
 175; notice of his memoir on the
 osteology of *Numenius longiro-
 stris*, 208; on the feeding habits of
 Phalacroptila nuttalli, 382; an-
 nouncement of his work entitled
 'Contributions to the Anatomy
 of North American Vertebrates,'
 391.
Sialia arctica, 8, 15, 140, 234.
 mexicana, 15, 140, 172, 331,
 349.
 sialis, 15, 140, 277, 381.
 sialis azurea, 85.
Simorhynchus cristatellus, 159, 341,
 388.
 psittacula, 159, 342.
 pygmæus, 159.
Sitta canadensis, 15, 277.
 aculeatus, 15, 126, 172, 277,
 331, 349.
 carolinensis, 277, 321, 349.
 pygmæa, 15, 126, 172, 331,
 350.
Siurus auricapillus, 15, 61, 62, 65,
 133, 278.
 motacilla, 91.
 nævius, 15, 278.
Skimmer, Black, 193.
Skylark, Missouri, 369.
Smith, Hugh M., breeding of *Loxia*
 americana in the District of Co-
 lumbia 379; *Ereunetes occidenta-
 lis* on the lower Potomac, 385;
 Harelda hyemalis in Maryland in
 summer, 385.
Snipe, American, 336.
 Red-bellied, 37, 286.
 Red-breasted, 374.
 Wilson's 286.
Snowbird, 115, 117.
 Black, 98, 265, 280.
 White-winged, 32.
Solitaire, Townsend's, 128.
Somateria dresseri, 111, 113.
 mollissima, 111.
 spectabilis, 63.
 v-nigra, 63, 158, 386.
Southwick & Jencks, southern range
 of *Rissa tridactyla kotzebuei*, 313.
Sparrow, Baird's, 198, 333.
 Clay-colored, 280.
 Chipping, 145, 280.
 English, 190, 211, 296.
 Field, 145, 280.
 Fox-colored, 150, 281.
 House, 99.
 Harris's, 281.
 Henslow's, 190.

- Sparrow, Ipswich, 36, 97, 99, 105.
 Java, 314.
 Large-billed, 198.
 Leconte's, 23, 190, 280, 381.
 Lincoln's, 280.
 Oak-woods, 299.
 Savanna, 97, 272, 280.
 Song, 95, 147, 236, 263, 280.
 Swamp, 100, 149, 189, 216, 280, 307, 380.
 Titlark, 212.
 Tree, 144, 236, 262, 265, 280.
 White-crowned, 272, 281.
 White-throated, 281.
 Worthen's Field, 291.
 Yellow-winged, 280.
- Spatula clypeata, 18, 112, 193, 288.
- Speotyto cunicularia hypogæa, 9, 17, 143, 284.
- Spermophila, 47.
- Sphyrapicus thyroides, 17, 174, 349.
 varius, 17, 258.
 varius nuchalis, 9, 17, 174.
- Spindalis benedicti, 295.
 exsul, 295.
- Spiza americana, 106, 192, 281.
- Spizella breweri, 16.
 domestica, 145, 280.
 domestica arizonæ, 16, 141.
 montana, 16.
 monticola, 144, 280.
 monticola ochracea, 236.
 pallida, 16, 141, 280.
 pusilla, 145, 280.
 socialis, 380.
 wortheni, 291.
- Sporophila, 47.
 moreletii, 47.
- Squatarola helvetica, 32, 63, 285.
- Steganopus wilsoni, 18, 273, 286.
- Stejneger, Leonard, *Analecta ornithologica*: XVII, on the origin of the word *Quiscalus*, 43; XVIII, *Colinus*, not *Ortyx*, 44; XIX, is the name *Ortyx massena* tenable? 46; XX, *Cyanolesbia*, strictly correct, 46; XXI, concerning the names of some North American Spizinae, 47; XXII, the correct name of the American Cowbird, 49; XXIII, remarks on the generic name *Sayornis* and on *Sayornis phæbe*, 51; XXIV, on the proper generic name of the Pileated Woodpecker and allies, 51; XXV, Why *Chordeiles virginianus* and not *Ch. popetue*? 178; XXVI, on the generic names of the Phalaropes, 182; XXVII, *Scops* preoccupied, 183; XXVIII, on Gyrfalcons, 184; XXIX, *Facana* contra *Parra*, 337; *Colymbus nigricollis*, not a North American Bird, 340; XXXI, "*Simorhynchus* Merrem, 1819," 341; XXXII, *Dendroica vigorsii* (Aud.), 343; notice of his paper on the Wrens of the subgenus *Anorthura*, 94; do. the Ptarmigans of the group *Attagen*, 94; do. descriptions of new species of birds from Kamtschatka, 95; do. list of recent ornithological publications in the United States, 95; notice of his paper on the genus *Cephus*, 294; on the alleged occurrence of the Pacific Eider in Labrador, 386.
- Stelgidopteryx serripennis, 15, 141, 279.
- Stephens, F., notes of an ornithological trip in Arizona and Sonora, 225.
- Stercorarius crepidatus, 63.
 longicaudus, 158.
 parasiticus, 18, 63, 158, 232.
 pomatorhinus, 63.
- Sterna anæsthesia, 230.
 anglica, 38.
 antillarum, 289.
 forsteri, 18, 289.
 macrura, 63.
 paradisea, 158.
- Streptopelia interpres, 63.
- Strix nebulosa, 189, 284.
- Sturnella magna, 250.
 magna mexicana, 198.
 neglecta, 13, 16, 142, 282.
- Sula bassana, 98.
- Surnia funerea, 31, 108, 220.
 ulula caparoch, 383.
- Swallow, Bank, 87, 279.
 Barn, 278, 266.
 Cliff, 279, 366.
 European Chimney, 366.
 Rough-winged, 279.
 Violet-green, 279, 333.
 White-bellied, 86, 98, 189, 278.
- Swan, Trumpeter, 287.
 Whistling, 10, 193.
- Swift, Chimney, 257, 283.
 White-throated, 9.
- Sylvania microcephala, 372.
- Sylvia pinus, 343.
 solitarius, 343.
 vigorsii, 343.
- Symphemia semipalmata, 18, 286, 289.

- Synthliborhamphus antiquus, 159.
 TACHYCINETA bicolor, 15, 278.
 thalassina, 15, 141, 279, 333, 354.
 Tachypetes aquila, 113.
 Tanager, Cooper's 198.
 Liver-colored, 197.
 Louisiana, 333.
 Scarlet, 278.
 Western, 8.
 Tantalus loculator, 18, 221, 311.
 Teal, Blue-winged, 112, 288, 375.
 Green-winged, 193, 238, 288, 295.
 Telmatodytes palustris, 15, 190, 278.
 Tell-tale, Greater, 192.
 Lesser, 189.
 Teretrictis fernandina, 368.
 fornsi, 368.
 Tern, Black, 112.
 Gull-billed, 38.
 Thalassidroma pelagica, 99.
 Thompson, Frank J., recent occurrence of the Black Vulture in Ohio, 109; the *Ædicnemus dominicensis* in confinement, 110.
 Thrasher, Brown, 277.
 Leconte's, 196.
 Palmer's, 196.
 Sage, 8.
 Thrush, Brown, 98, 277, 298.
 Dwarf, 330.
 Golden-crowned, 98, 278.
 Long-billed Water, 91.
 Olive-backed, 98, 277.
 Rocky Mountain Hermit, 330.
 Russet-backed, 85.
 Small-billed Water, 278.
 Swainson's, 330.
 Wilson's, 277.
 Wood, 72, 215, 277, 288.
 Thryomanes bewicki leucogaster, 140.
 Titlark, 209, 278.
 Tinnunculus sparverius, 17, 143, 285.
 sparverius isabellinus, 205.
 Titmouse, Tufted, 277.
 Totanus flavipes, 18, 143, 189, 286.
 melanoleucus, 18, 99, 193, 286.
 solitarius, 374.
 Towhee, 282.
 Cañon, 237.
 Northern, 282.
 Spurred, 128.
 Townsend, Charles H., the occurrence of the Catbird (*Mimus carolinensis*) on the Farallone Islands, Pacific Ocean, 215.
 Townsend, Charles W., the Black-throated Bunting in Maine, 106.
 Tringa canutus, 63, 312, 375.
 maculata, 157, 374.
 minutilla, 376.
 Tringoides macularius, 18, 143, 286, 298.
 Trochilus alexandri, 310.
 colubris, 17, 218, 256, 283, 299.
 Troglodytes aëdon, 140, 278.
 aëdon marianæ, 351.
 aëdon parkmani, 15, 278.
 alascensis, 94, 157.
 beani, 295.
 borealis, 94.
 fumigatus, 94.
 hiemalis, 94.
 hiemalis pacificus, 94.
 parvulus, 94.
 parvulus bergensis, 94.
 Tryngites rufescens, 63, 286.
 Turdus migratorius, 102, 114.
 mustelinus, 215.
 unalascæ auduboni, 349.
 ustulatus, 85.
 ustulatus aliciae, 190.
 Turkey, Wild, 10, 174, 285.
 Turner, Lucien M., notes on the birds of the Nearer Islands, Alaska, 154; notice of his 'List of the Birds of Labrador,' 368; familiar Chipping Sparrows, 380.
 Tyrannula cooperi, 208.
 mexicana, 208.
 Tyrannus carolinensis, 17, 142, 253, 283.
 tyrannus, 369.
 verticalis, 17, 143, 283.
 vociferans, 17.
 URIA grylle, 63.
 lomvia, 159.
 Urinator lumme, 159.
 pacificus, 159.
 Utamania torda, 38.
 VIREO approximans, 291.
 atricapillus, 274.
 bairdi, 295.
 belli, 101.
 carmioli, 291.
 gilvus, 279.
 gilvus swainsoni, 332, 354.
 huttoni stephensi, 197, 354.
 magister, 295.
 noveboracensis, 369.
 olivaceus, 65, 279.
 philadelphicus, 65, 272, 279, 305.
 pusillus, 197.

- Vireo solitarius*, 65, 279.
solitarius cassini, 332.
solitarius plumbeus, 332.
vicinior, 321.
Vireo, 117.
 Arizona, 321.
 Bell's, 101.
 Blue-headed, 279.
 Cassin's, 332.
 Gray, 321.
 Philadelphia, 272, 279, 305.
 Plumbeous, 332.
 Red-eyed, 98, 279.
 Stephens's, 197.
 Swainson's, 332.
 Warbling, 279.
 White-eyed, 72.
Vireolanus pulchellus verticalis, 293.
 cinerea, 295.
Vireosylva cinerea, 295.
 gilva, 141.
 grandior, 291.
Vultur californianus, 169.
Vulture, Black, 109.
 California, 167.
WARBLER, Audubon's, 329, 331.
 Bay-breasted, 272.
 Black-and-Yellow, 278.
 Blackburnian, 103.
 Black-capped, 278.
 Black-capped Yellow, 278.
 Black-poll, 278.
 Black-throated Gray, 197.
 Black-throated Green, 98.
 Blue-winged Yellow, 98, 378.
 Blue Yellow-backed, 97, 377.
 Cape May, 32, 33.
 Cerulean, 278.
 Connecticut, 104, 308.
 Golden Swamp, 32.
 Golden-winged, 102, 378.
 Grace's, 331.
 Hooded, 72, 190.
 Kirtland's, 376.
 Macgillivray's, 332.
 Mourning, 278.
 Olive, 172.
 Orange-crowned, 278, 330.
 Prothonotary, 97.
 Prairie, 278.
 Pine-creeping, 278, 343, 381.
 Red-poll, 104.
 Swainson's, 54, 65, 104, 105, 301, 305, 346, 377.
 Tennessee, 190.
 Townsend's, 331.
 White-throated, 378.
 Worm-eating, 103, 305.
Warbler, Yellow, 331.
 Yellow-bellied Red-poll, 190.
 Yellow-rumped, 98, 216, 278, 307, 380, 381.
 Yellow-throated, 72.
Waxwing, Cedar, 279.
 Northern, 279.
Widmann, Otto, note on the capture of *Coturniculus lecontei* and *Dendroæca kirtlandi* within the city limits of St. Louis, Mo., 381.
Wilcox, Timothy E., introduced game birds in Oregon and Idaho, 315.
Willard, Samuel Wells, food of the Hummingbird (*Trochilus colubris*), 218; the occurrence of *Chroicocephalus franklini* in Wisconsin, 222; notice of his paper on the birds of Brown and Outagamie Counties, Wis., 289.
Willet, 286, 298.
Wheatear, 113, 305.
Whip-poor-will, 179, 257, 272.
Wood, Dr. William, notice of death of, 391.
Woodcock, 99, 213, 261, 286.
Woodpecker, Black-backed, 265.
 Brown-headed, 174.
 California, 9, 174.
 Downy, 101, 257, 265, 283.
 Gairdner's, 126, 127.
 Golden-winged, 259, 284, 380.
 Hairy, 100, 265.
 Harris's, 9, 127, 174, 283.
 Nuchal, 174.
 Nuttall's, 213.
 Pileated, 51, 101, 283.
 Red-headed, 93, 99, 101, 209, 258, 283.
 Red-bellied, 283, 335.
 Red-naped, 9.
 Red-shafted, 9, 174.
 Strickland's, 199.
 Yellow-bellied, 258, 301.
Wren, Bewick's, 99.
 Cañon, 233.
 Carolina, 72, 86, 98, 209.
 House, 98, 278.
 Long-billed Marsh, 96, 190, 278.
 Rock, 234, 277.
 Short-billed Marsh, 190, 278.
 Western House, 278.
 Winter, 213, 115.
XANTHOCEPHALUS icterocephalus, 8, 16, 112, 192, 282.
Xema sabinei, 63.
YELLOWBIRD, Summer, 278.

- Yellow-legs, 286.
 Greater, 99, 286.
Yellow-shanks, 189.
Yellow-throat, Maryland, 72, 98,
 278, 367.
ZAMELODIA ludoviciana, 151, 192,
 281.
 melanocephala, 16, 99, 112,
 142, 281
Zeledon, José C., notice of his Cat-
 alogue of the Birds of Costa Rica,
 370.
Zenaida bogotensis, 358.
Zenaida rubripes, 357.
 ruficauda, 291, 357.
 vinaceo-rufa, 291, 359.
Zenaidura carolinensis, 9, 17, 143,
 285, 372.
 macroura, 372.
Zonotrichia albicollis, 65, 281.
 gambeli intermedia, 16, 141.
 intermedia, 124.
 leucophrys, 16, 65, 141, 272,
 281.
 querula, 281.